



Recombinant full-length tissue factor pathway inhibitor fails to bind to the cell surface: implications for catabolism in vitro and in vivo

Guyu Ho, Masaaki Narita, George J. Broze Jr, and Alan L. Schwartz

Tissue factor pathway inhibitor (TFPI) plays a key role in the regulation of tissue factor-initiated blood coagulation secondary to loss of the integrity of the blood vessel wall. TFPI is a naturally occurring Kunitz-type protease inhibitor that inhibits coagulation factor Xa and, in a factor Xa-dependent manner, mediates feedback inhibition of the factor VIIa/tissue factor

catalytic complex. In vivo full-length TFPI is thought to be primarily bound to the vascular endothelium and the high affinity binding requires an intact carboxy terminus. Here we describe a full-length TFPI molecule, expressed in mouse C127 cells (TFPI^{C127}), which exhibits virtually no cellular binding yet contains the intact carboxy terminus. This TFPI (TFPI^{C127}) is

neither internalized nor degraded via the TFPI endocytic receptor, LDL-receptor-related protein. Pharmacokinetic studies of TFPI^{C127} in vivo demonstrate a 10-fold prolongation in the plasma half-life, compared with that of bacterial recombinant TFPI. (Blood. 2000;95:1973-1978)

© 2000 by The American Society of Hematology

Introduction

The extrinsic pathway of coagulation is initiated by the exposure of tissue factor (TF) to the circulating blood.¹ Binding of factor VII to TF promotes the activation of factor VII to factor VIIa. The factor VIIa/TF complex activates its substrate factor IX and factor X, triggering a cascade of events that culminates in the formation of a fibrin clot.^{2,3} Tissue factor pathway inhibitor (TFPI), a 42 kd plasma glycoprotein, plays a key role in the regulation of TF-initiated blood coagulation^{2,3} via its abilities to directly inhibit factor Xa and, in a factor Xa dependent manner, inhibit the factor VIIa-TF proteolytic activity.³ Mature TFPI consists of a negatively charged amino terminus, 3 tandem Kunitz-type inhibitory domains, and a positively charged carboxy terminus.⁴ The second Kunitz domain binds and inhibits factor Xa, while the first Kunitz domain binds and inhibits the factor VIIa/TF complex.⁴ The carboxy terminus of TFPI and a portion of the third Kunitz domain contain heparin-binding sites^{5,6} and are necessary for the anticoagulant function of TFPI in TF-induced coagulation in vitro⁵ and for TFPI binding to the cell surface.⁷

TFPI circulates in plasma predominantly bound to lipoproteins.⁸ Platelets contain approximately 8% of the total TFPI in blood and release their TFPI after stimulation with thrombin.⁹ The major endogenous source of TFPI is thought to be bound to the vascular endothelium and is releasable after heparin infusion, whereupon plasma TFPI levels rise several fold.^{10,11}

Recent animal studies have demonstrated that recombinant TFPI is effective against TF-induced coagulopathy,¹² prevents arterial thrombosis,^{13,14} and reduces mortality from bacterial septic shock.¹⁵ Pharmacokinetic studies after an intravenous bolus injection of recombinant TFPI have shown that TFPI is rapidly cleared from the circulation with a plasma half-life of approximately 2 minutes in rabbits¹⁶ and less than 1 minute in rats.⁷ As a result, high doses of recombinant TFPI (approximately 20 mg/kg/d) are required to achieve therapeutic efficacy.¹⁷

We have recently shown that the rapid clearance of recombinant TFPI from the circulation is in large part dependent on the binding of TFPI to the vascular endothelium and to a lesser degree to its hepatic removal via the endocytic receptor LDL-receptor-related protein (LRP).¹⁸ In this study, we demonstrate that full-length TFPI expressed in mouse C127 cells does not bind to the cell surface nor is it degraded via LRP in vitro. This results in an approximately 10-fold prolongation of the TFPI^{C127} plasma half-life in mice. The noncellular binding properties of TFPI^{C127} may thus offer additional therapeutic potential.

Materials and methods

Materials

Iodogen was purchased from Pierce (Rockville, IL). [¹²⁵I]iodide was from Amersham Corp (Piscataway, NJ). Human factor Xa was from Enzyme Research (South Bend, IN) and Spectrozyme Xa was from American Diagnostica, Inc (Greenwich, CT). Bovine serum albumin was purchased from Calbiochem-Novabiochem. Affi-gel 10 was from Bio-rad (Hercules, CA). Tissue culture media and plasticware were obtained from Life Technologies, Inc (Rockville, MD). Recombinant TFPI from *Escherichia coli* was provided by Monsanto Company (St Louis, MO). CHO, SK hepatoma, and 293 cell-derived TFPI, was purified as described below.

Purification of TFPI^{C127}.—TFPI^{C127} was affinity-purified from the conditioned media of mouse C127 cells that had been transfected with a bovine papilloma virus containing wild-type human TFPI cDNA.¹² In brief, cells producing TFPI^{C127} were cultured in serum-free media containing apolipoprotein (10 µg/mL; Sigma) and the media were replaced every 2 to 3 days. Harvested media were concentrated using an Amicon ultraconcentrator (YM 30; Amicon, Bedford, MA) and the concentrates were absorbed to a 2F8 monoclonal anti-TFPI-affigel 10 column (0.5 × 3 cm) and eluted with 100 mmol/L glycine, pH 2.2. The pH of the eluant was adjusted to pH 7.5 using 1 mol/L Tris-HCl, pH 8.0, and applied to a 1 mL HiTrap

From the Departments of Pediatrics, Molecular Biology and Pharmacology, Medicine, and Cell Biology and Physiology, Washington University School of Medicine and St Louis Children's Hospital, St Louis, MO

Submitted June 15, 1999; accepted November 2, 1999.

Supported by grants from the NIH and Monsanto/Searle.

Reprints: Alan L. Schwartz, Department of Pediatrics, Campus Box 8116.

Washington University School of Medicine, 660 S Euclid Ave, St Louis, MO 63110.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 U.S.C. section 1734.

© 2000 by The American Society of Hematology

heparin-agarose column (Pharmacia-LKB Biotechnology, Piscataway, NJ), which had been equilibrated in 0.1 mol/L NaCl, 0.05 mol/L Tris-HCl, pH 7.5. The column was developed with a linear NaCl gradient from 0.1 to 1.0 mol/L at 0.5 mL/min over 10 column volumes. Fractions were assayed for the presence of TFPI^{C127} by a modified enzyme linking immunosorbent assay (ELISA) method using 2 independent monoclonal antibodies, 2H8 and 2B12.¹ The desired fractions were pooled, concentrated using Centricon microconcentrators (Amicon), and stored at -80°C. The protein concentration was determined using the modified ELISA method with TFPI of known concentration as standard.

Amidolytic assays of factor Xa activity

In a 96-well microtiter plate, 1 nmol/L of factor Xa was incubated with various concentrations of TFPI (0-3 nmol/L) in 135 μ L of 0.1 mol/L NaCl, 0.05 mol/L Tris-HCl, 0.5% bovine serum albumin (BSA), pH 7.5 (TBSA) at room temperature for 30 minutes. Fifteen microliters of 1 nmol/L Spectrozyme Xa chromogenic substrate was then introduced (final concentration 100 μ mol/L) and the change in absorbance at 405 nm was determined 10 minutes thereafter.

Tissue factor pathway inhibitor functional assay of VIIa/TF inhibition

The end point assay of anti-VIIa/TF activity (American Diagnostica) was used to compare TFPI^{C127} and bacterially expressed TFPI at equimolar concentrations.

Factor Xa-induced coagulation of plasma

In a fibrometer (BBL, Cockeysville, MD), 50 μ L of rabbit brain cephalin prepared as described by the manufacturer (Sigma), 50 μ L of CaCl₂ (25 mmol/L), 50 μ L of various amounts of TFPI^{C127} (25 ng to 300 ng) in TBSA, and 50 μ L of human factor Xa (0.2 nmol/L) were incubated at 37°C. After 30 seconds, 50 μ L of normal human plasma (George King Biochemical) were added, and the degree of apparent factor Xa inhibition was determined by comparing the clotting to a factor Xa standard curve.

Cell culture

Human hepatoma HepG2 cells, mouse fibroblast P8A 13 cells,¹⁹ and mouse brain endothelial cells (bend-3)²⁰ were cultured in Dulbecco's modified Eagle's medium (DMEM), supplemented with 10% fetal calf serum, 100 U/mL of penicillin, and 100 μ g/mL of streptomycin. Cells were incubated at 37°C in humidified air containing 5% CO₂.

Protein iodination

Proteins (5-25 μ g) were iodinated using the IODOGEN method.²¹ Specific radioactivities were typically 1 to 10 $\times 10^6$ cpm/pmol of protein.

Binding and degradation assays

Binding of ¹²⁵I-TFPI to cells was performed in suspension. Cells were displaced from Petri-dishes by incubation with 3 mmol/L EDTA in phosphate-buffered saline (PBS) at room temperature for 5 minutes. Cells were then pelleted at low speed (1000 rpm \times 5 minutes) on a table top centrifuge (Sorvall RT6000B) to remove EDTA and resuspended in the assay buffer (DMEM/3% BSA). 4×10^6 cells in a volume of 300 μ L were added with 2 nmol/L of ¹²⁵I-TFPI in the presence or absence of 100 mol excess of cold ligands. After rocking at 4°C for 2 hours, cell suspensions were gently layered on 900 μ L of fetal calf serum in microfuge tubes and spun at 14 000 rpm \times 1 minute in a table-top microfuge. The cell pellet was counted in a gamma counter (Cobra II auto-gamma, Packard Instrument, Meriden, CT). Nonspecific binding was determined in the presence of excess unlabeled ligand as specified in the figure legends.

Degradation assays were performed by washing cell monolayers (in 12-well dishes) twice with the assay buffer. 0.5 mL of assay buffer containing 2 nmol/L ¹²⁵I-TFPI was then added to each well. After incubation at 37°C for the indicated periods, the overlying medium was removed and proteins were precipitated by the addition of BSA to 5 mg/mL

and trichloroacetic acid to 20%. Degradation of ligand was defined as the appearance of radioactive fragments in the overlying medium that were soluble in trichloroacetic acid. Fragmentation of ¹²⁵I-TFPI at 37°C in cell-free wells was subtracted from the corresponding samples.

For ligand competition binding and degradation assays (Table 2), HepG2 and CHO cells were assayed as monolayers as described previously.¹⁴ ¹²⁵I-TFPI was at 20 nmol/L for the binding assays and at 0.6 nmol/L for the degradation assays.

Plasma clearance of ¹²⁵I tissue factor pathway inhibitors in mice

Twelve- to 16-week-old BALB/c mice (weighing 20-25 g) were anesthetized with sodium pentobarbital (1 mg/20 g of body weight) during the course of experiments. Approximately 15 pmol of radiolabeled protein alone or with unlabeled 39 kd protein in 100 μ L of sterile saline was injected into the tail vein over 30 seconds. In studies in which heparin was administered, heparin was injected 20 minutes after ¹²⁵I-TFPI administration. At the indicated times, 40 to 50 μ L of blood was collected by periorbital bleeding. The blood samples were centrifuged and 15 μ L of the plasma fractions was spotted onto 3 MM filter paper (Whatman Inc, Clifton, NJ), precipitated with 10% trichloroacetic acid, rinsed with ethanol, and radioactivity determined. The initial plasma concentration (time = 0) of ¹²⁵I-TFPI was extrapolated from the measured points. At the end of each experiment, animals were killed and the liver, kidneys, spleen, and lungs were removed, blotted, weighed, and the radioactivity determined.

Metabolic incorporation of ³⁵SO₄ and immunoprecipitation of TFPI^{C127}

One 75-cm² flask of C127 cells was labeled for 96 hours with 37 MBq (1 mCi) of Na₂³⁵SO₄ (Amersham Corp) in 10 mL of sulfate-deficient DMEM (-SO₄) supplemented with 2% dialyzed fetal calf serum, in the absence of G418. The culture medium was then concentrated using Centricon microconcentrators (Amicon). Monoclonal antibody 2H8-cojugated affigel-10 beads were added to the concentrates that were then rocked overnight to allow for binding. The beads were subsequently collected by centrifugation, washed 3 times with PBS, boiled in SDS-sample buffer (to release bound TFPI), and the bead-free portion was analyzed by SDS-PAGE.

Electrophoresis and Western blotting

SDS-PAGE was performed using 10% separating and 5% stacking gels and proteins were transferred to nitrocellulose membrane. The membrane was blocked with 5% nonfat milk for 30 minutes, followed by incubation with a 1:100 dilution of rabbit antisera raised against a synthetic peptide matching the carboxy terminal 12 amino acid residues of the mouse TFPI protein.⁵ The immunoreactivity was detected using ECL reagents (Amersham Corp).

Results

Purification of recombinant tissue factor pathway inhibitor from mouse C127 cells

Previous studies have shown that TFPI, when expressed in mammalian systems, is susceptible to proteolysis at its carboxy terminus.^{5,22} Thus, mouse C127 cells expressing TFPI were grown in media containing the protease inhibitor, aprotinin. TFPI was purified from the media via TFPI monoclonal antibody-affinity chromatography. This was followed by heparin-agarose affinity which separates the full-length TFPI from the truncated species by charge interactions between the heparin and the positively charged carboxy terminus of TFPI. Figure 1 shows the elution of TFPI^{C127} from the heparin column with a linear NaCl gradient. TFPI^{C127}




Figure 1. Profile of heparin-agarose chromatography of TFPI^{C127}. Heparin-agarose chromatography was carried out using a 0.1 to 1 mol/L NaCl gradient in 0.05 mol/L Tris-HCl, pH 7.5. Elution of TFPI^{C127} was monitored by a modified ELISA as described in the "Materials and Methods." Inset: Western blot analysis of TFPI^{C127}. Purified TFPI^{C127} (1 µg) was run on a 10% SDS-PAGE gel and transferred to nitrocellulose membrane. The membrane was blotted with rabbit antiserum containing a peptide antibody against the last 12 amino acid residues of TFPI, and developed with ECL reagents.

eluted with 2 peaks at NaCl concentrations of 0.4 and 0.6 mol/L, respectively. Because full-length TFPI elutes at 0.6 mol/L salt,^{5,6} only the fractions collected at or greater than 0.6 mol/L NaCl (fractions 12-15) were pooled for subsequent studies. To confirm that the purified TFPI^{C127} was full length, TFPI^{C127} was subjected to Western blot analysis using antibodies directed against the last 12 amino acid residues of TFPI. As shown in the inset of Figure 1, the TFPI was recognized by these antibodies, confirming that the carboxy terminus was intact.

The activity of TFPI^{C127} was assessed in a direct amidolytic assay of factor Xa activity, as well as an end point assay of




Figure 2. Inhibition of factor Xa amidolytic activity by TFPI^{C127}. Factor Xa (1 nmol) was incubated with increasing concentrations of TFPI^{C127} (0-3 nmol) at room temperature for 30 minutes. The remaining factor Xa activity was assessed using the chromogenic substrate Spectrozyme Xa. Bacterial recombinant TFPI (bTFPI) was assayed in the identical manner and used as control.

Table 1. Binding of ¹²⁵I-TFPI^{C127} to various cell lines

Cell Line	Specific Binding (fmol/10 ⁶ cells)	
	¹²⁵ I-TFPI ^{C127}	¹²⁵ I-bTFPI
HepG2	Not detectable	580
Bend	Not detectable	810
PEA 13	Not detectable	540

HepG2, Bend, and PEA 13 cells were incubated in suspension with 2 nmol ¹²⁵I-TFPI^{C127} in the presence or absence of 100 mol excess of cold TFPI^{C127} for 2 hours at 4°C. The specific radioactivity associated with cells was determined (open bars). Binding of bacterial recombinant ¹²⁵I-TFPI to these cells was performed in the identical manner and used as control. Values represent the means of duplicate determinations.

anti-VIIA/TF activity. As shown in Figure 2, TFPI^{C127} was able to inhibit factor Xa activity in a concentration-dependent manner with complete inhibition of factor Xa at approximately 3 nmol/L, which was essentially identical to that of bacterial recombinant TFPI. Using a global anticoagulant assay, TFPI^{C127} inhibited factor Xa-induced plasma coagulation with IC₅₀ approximately 10 nmol/L, compared with approximately 4 nmol/L for the bacterial recombinant TFPI. The disparity in IC₅₀ values observed here is consistent with the previous observations that TFPI expressed in prokaryotic systems exhibits higher anticoagulant activities than that expressed in eukaryotic systems. Using the end point assay that measures anti-VIIA/TF activity, on a molar the activity of TFPI^{C127} is 108% ± 16% that of bacterially expressed TFPI.

Cellular binding and degradation of tissue factor pathway inhibitor in C127 cells

We have previously shown that bacterial recombinant TFPI binds to human hepatoma HepG2 cells with high affinity²⁴ and that truncation of sequences distal to the second Kunitz domain abolishes the binding.⁷ Surprisingly, the full-length ¹²⁵I-TFPI^{C127} exhibited no binding to hepatoma HepG2 cells, or to mouse vascular endothelial Bend cells, or mouse fibroblast PEA 13 cells (Table 1). Bacterial recombinant ¹²⁵I-TFPI, however, exhibited abundant binding to these cells (Table 1). Furthermore, ¹²⁵I-TFPI^{C127} was not taken up/degraded by HepG2 cells in contrast to




Figure 3. Degradation of ¹²⁵I-TFPI^{C127} by HepG2 cells. HepG2 cells growing in petri dishes were incubated with 0.5 nmol of ¹²⁵I-TFPI^{C127} at 37°C for the indicated times. At each time point, the overlying medium was removed and proteins were precipitated by trichloroacetic acid (TCA). The TCA-soluble radioactivity (ie, degradation products) was determined. Degradation of bacterial recombinant ¹²⁵I-TFPI by HepG2 cells was performed in the identical manner and served as a control. Numbers represent the means of duplicate determinations. Total degraded ¹²⁵I-TFPI at 4 hours was less than 10% input.




Figure 5. $^{35}\text{SO}_4$ is incorporated into TFPI^{C127}. C127 cells expressing human TFPI were incubated with $^{35}\text{SO}_4$ for 85 hours, after which the culture medium was collected and immunoprecipitated with anti-TFPI as described under "Materials and Methods." The immunoprecipitates were analyzed via SDS-PAGE and autoradiography (left), or via Western blot analysis using TFPI monoclonal antibody 24-8 (right).

Binding of full-length TFPI to the cell surface is dependent on its highly positively charged carboxyl terminus that interacts with the negatively charged cell surface HSPGs.^{7,10,28} We have previously shown that TFPI expressed in bacteria binds with high affinity to heparoma HepG2 cells, fibroblast PEA13 cells, and microvascular endothelial cells.^{20,24,27} Similarly, TFPI expressed in hepatoma and CHO cells displays similar binding properties (Table 3). To our surprise, TFPI expressed in mouse C127 fibroblast cells was devoid of cell surface binding activity but remained biologically active as judged by the inhibition of (1) factor Xa amidolytic activity (Figure 2), of (2) factor Xa-induced plasma coagulation, and (3) factor VIIIa/TF activity. One potential explanation for this loss of binding activity is that secondary to intramolecular interactions, TFPI's carboxyl terminus is no longer available for interaction with cell surface HSPGs. It is of note that, although TFPI^{C127} does not bind to cell surface HSPGs, it binds to heparin-agarose (Figure 1) with the same affinity as bacterial recombinant TFPI, ie, both TFPI species are eluted from heparin-agarose at identical salt concentrations (0.6 mol/L NaCl) (6, data not shown). The fact that the affinity of TFPI is 40-fold greater for heparin than for HSPGs²⁸ may account for the high affinity binding of both species of TFPI to heparin and their discordant binding to cell surface HSPGs.

Girard et al²⁹ have shown that serine 2 of TFPI is phosphorylated after its expression in C127 cells. However, TFPI with a serine 2 to alanine mutation, when expressed in these cells, did not regain the binding properties of wild type TFPI (data not shown).

This finding suggests that this negatively charged phosphate group (ie, at serine 2) does not interfere with availability of the carboxyl terminus. TFPI contains 3 potential N-linked and 1 O-linked glycosylation sites.³⁰ Smith et al²⁴ have shown that metabolic labeling of 293 cells with [^{35}S]SO₄ yielded sulfated TFPI, whereas sulfated TFPI was not produced by CHO cells. They further determined that TFPI²⁹³ carries Asn-linked sulfated oligosaccharides: SO₄-GalNAc β 1,2GlcNAc β 1,2Man α .²⁶ Because TFPI²⁹³ was unable to bind to the cell surface (Table 3), it is possible that sulfated oligosaccharides on TFPI may interact with its own positively charged carboxyl terminus, thus abrogating TFPI's interaction with the cell surface. Attempts to quantitatively remove linked oligosaccharides or sulfate via incubation of nondenatured TFPI^{C127} with N-glycosase, O-glycosase or sulfatase and assess biological function have proven unsuccessful. Thus, whether TFPI^{C127} possesses the same sulfation pattern as TFPI²⁹³ will require further study.

TFPI^{C127} exhibited a 10-fold increase in plasma half-life compared with bacterial recombinant TFPI (Figure 4A). Because the administration of heparin did not significantly increase the plasma [^{125}I]-TFPI^{C127} concentration, secondary to release from the vascular endothelium, or did the administration of the 39-kD protein inhibit its clearance (Figure 4), the prolonged half-life of [^{125}I]-TFPI^{C127} is likely due to its inability to bind to either cell surface HSPGs or LRP.³¹ These results are consistent with our *in vitro* findings (Figure 3 and Table 1). However, it is not clear at present whether this observation can be accounted for by alterations in protein sulfation alone, because [^{125}I]-TFPI²⁹³ and [^{125}I]-TFPI^{C80} have similar rates of plasma clearance.²⁶ Nonetheless, the clearance of [^{125}I]-TFPI^{C127}, albeit markedly prolonged, did result in elimination of 50% at approximately 20 minutes. Therefore, additional mechanisms (eg, hepatic glycoprotein receptors) may be operative.

Under physiologic conditions, TFPI is primarily synthesized and secreted by vascular endothelial cells.³² Because the major pool of TFPI *in vivo* is associated with the endothelial cell surface, it appears that endothelial TFPI maintains the carboxy terminus available for interaction with cell surface HSPGs. Assuming that sulfation does play a role in TFPI binding to cells, TFPI produced by endothelial cells *in vivo* may well have to be under- or unsulfated to maintain its cellular binding. However, it is not presently known whether endogenous TFPI is sulfated or its degree of sulfation. Of note, human endothelial cells express significantly lower (less than 10%) GalNAc- and sulfotransferase activities than those expressed in 293 cells. These 2 enzymes are the key activities required for the addition of sulfated oligosaccharides to TFPI.²⁶

The physiologic significance of the cell-associated TFPI pool may relate to both its rapid mobilization into the plasma as well as its role as a mediator in the cellular uptake and degradation of factor Xa.²⁷ In addition, it may serve to down-regulate the coagulant activity of factor VIIIa/TF complex via transfer to caveolae.³³ Endogenous TFPI within the circulation, however, is likely of less physiological importance, as it is largely truncated at various positions within the carboxy terminal domain³⁴ and hence exhibits much reduced anticoagulant activity.³ The nonbinding full-length TFPI we have characterized in this study may therefore offer a therapeutic advantage when high plasma TFPI concentrations are required to effectively modulate TF-induced blood coagulation.

References

- Wojcik MJ, Turitto VT, Baumgartner HR, Hemerson Y, Hoffman T. Evidence for the presence of tissue factor activity on endothelium. *Blood*. 1989;73:368.
- Rapaport SI. Inhibition of factor VIIa/tissue factor-induced blood coagulation with particular emphasis upon a factor Xa-dependent inhibitory mechanism. *Blood*. 1989;73:358.
- Broze GJ Jr. The role of tissue factor pathway inhibitor in a revised coagulation cascade. *Scand Hematol*. 1992;29:158.
- Girard TJ, Warren LA, Novotny WF, et al. Functional significance of the Kunitz-type inhibitory domains of lipoprotein-associated coagulation inhibitor. *Nature*. 1989;336:516.
- Wesselschmidt R, Liberti KM, Girard TJ, Wan TC, Broze GJ Jr. Tissue factor pathway inhibitor: the carboxy-terminus is required for optimal inhibition of factor Xa. *Blood*. 1992;79:2004.
- Wesselschmidt R, Liberti KM, Huang ZF, MacPhail L, Broze GJ Jr. Structural requirements for tissue factor pathway inhibitor interactions with factor Xa and heparin. *Blood Coagul Fibrinolysis*. 1993;4:561.
- Wanawatsky I, Bu G, Mesi A, Saffitz JE, Broze GJ Jr, Schwartz AL. The carboxy terminus of tissue factor pathway inhibitor is required for interacting with heparan sulfate in vitro and in vivo. *J Clin Invest*. 1995;95:1773.
- Novotny WF, Girard TJ, Milotich JP, Broze GJ Jr. Purification and characterization of the lipoprotein-associated coagulation inhibitor from human plasma. *J Biol Chem*. 1989;264:18,532.
- Novotny WF, Girard TJ, Milotich JP, Broze GJ Jr. Platelets secrete a coagulation inhibitor functionally and antigenically similar to the lipoprotein-associated coagulation inhibitor. *Blood*. 1989;72:2020.
- Sandset PM, Abildgaard U, Larsen ML. Heparin induces release of extrinsic coagulation pathway inhibitor (EPI). *Thromb Res*. 1986;50:503.
- Novotny WF, Brown BG, Milotich JP, Broze GJ Jr. Plasma antigen levels of the lipoprotein-associated coagulation inhibitor in patient samples. *Blood*. 1991;78:387.
- Day KO, Hoffman LC, Palmer MO, et al. Recombinant lipoprotein-associated coagulation inhibitor inhibits tissue thromboplastin-induced intravascular coagulation in the rabbit. *Blood*. 1990;76:1538.
- Hastak EJ, Torr SR, Day KC, et al. Prevention of arterial reocclusion after thrombolysis with recombinant lipoprotein-associated coagulation inhibitor. *Circulation*. 1991;84:821.
- Undahl AK. Tissue factor pathway inhibitor: a potent inhibitor of in vitro coagulation and in vivo thrombus formation. *Curr Opin Lipidol*. 1994;5:434.
- Onasch AA, Chang ACK, Folson L, Wan TC, Taylor FB, Hershaw LB. Tissue factor pathway inhibitor reduces mortality from *Escherichia coli* septic shock. *J Clin Invest*. 1993;91:2350.
- Palmer MO, Hall LJ, Ralston CM, Baldwin MK, Wilson AGE, Wan TC. Clearance of recombinant tissue factor pathway inhibitor (TFPI) in rabbits. *Thromb Haemostas*. 1992;68:33.
- Girard TJ. Tissue factor pathway inhibitor. In: Sakahara A, Loscalzo J, eds. *Novel Therapeutic Agents in Thrombosis and Thrombolysis*. New York: Marcel Dekker; 1997:225.
- Nasir M, Bu G, Chen GM, et al. Two receptor systems are involved in the plasma clearance of tissue factor pathway inhibitor in vivo. *J Biol Chem*. 1995;270:24,800.
- Withrow TE, Herz J. Genetic deficiency in low density lipoprotein receptor-related protein confers cellular resistance to *Pseudomonas aeruginosa* A. Evidence that this protein is required for uptake and degradation of multiple ligands. *J Cell Sci*. 1994;107:719.
- Ho G, Broze GJ Jr, Schwartz AL. Role of heparan sulfate proteoglycans in the uptake and degradation of tissue factor pathway inhibitor-coagulation factor Xa complexes. *J Biol Chem*. 1997;272:18,538.
- Bu G, Morton PA, Schwartz AL. Identification and partial characterization by chemical cross-linking of a binding protein for tissue-type plasminogen activator (t-PA) on rat hepatoma cells. A plasminogen activator inhibitor type 1-independent t-PA receptor. *J Biol Chem*. 1992;267:15,585.
- Nordling O, Bjorn SE, Valentin S, et al. The C-terminus of tissue factor pathway inhibitor is essential to its anticoagulant activity. *Biochemistry*. 1991;30:10,371.
- Diaz-Coller JA, Palmer MO, Kreitzer KK, et al. Isolation and characterization of recombinant tissue factor pathway inhibitor expressed in *Escherichia coli*. *Thromb Haemostas*. 1994;71:538.
- Wanawatsky I, Broze GJ Jr, Schwartz AL. The low density lipoprotein receptor-related protein mediates the cellular degradation of tissue factor pathway inhibitor. *Proc Natl Acad Sci U S A*. 1994;91:9884.
- Bu G. Receptor-associated protein: a specialized chaperone and antagonist for members of the LDL receptor gene family. *Curr Opin Lipidol*. 1996;7:149.
- Smith PL, Skolton TP, Field D, et al. The asparagine-linked oligosaccharides on tissue factor pathway inhibitor terminate with SO4-4GalNAc beta 1,4GalNAc beta 1.2 Mann alpha. *J Biol Chem*. 1992;267:19,140.
- Ho G, Toomey JR, Broze GJ Jr, Schwartz AL. Receptor-mediated endocytosis of coagulation factor Xa requires cell surface-bound tissue factor pathway inhibitor. *J Biol Chem*. 1995;270:19497.
- Valentin S, Lammert A, Ostergaard P, Nielsen J, Nordling O. Characterization of the binding between tissue factor pathway inhibitor and glycosaminoglycans. *Thromb Res*. 1994;75:173.
- Girard TJ, McCourt D, Novotny WF, MacPhail L, Liberti KM, Broze GJ Jr. Endogenous phosphorylation of the lipoprotein-associated coagulation inhibitor at serine-2. *Biochem J*. 1990;270:621.
- Wan TC, Kreitzer KK, Girard TJ, Milotich JP, Broze GJ Jr. Cloning and characterization of a cDNA coding for the lipoprotein-associated coagulation inhibitor shows that it consists of three tandem Kunitz-type inhibitory domains. *J Biol Chem*. 1988;263:6001.
- Schwartz AL, Broze GJ Jr. Tissue factor pathway inhibitor endocytosis. *Trends Cardiovasc Med*. 1997;7:234.
- Bajaj MS, Kuppenswamy MN, Saito H, Spitzer SG, Bajaj SP. Cultured normal human hepatocytes do not synthesize lipoprotein-associated coagulation inhibitor: evidence that endothelium is the principal site of its synthesis. *Proc Natl Acad Sci U S A*. 1990;87:5868.
- Gevinsky JR, Rao SYA, Ruf W. Ligand-induced protease receptor translocation into caveolae: a mechanism for regulating cell surface proteolysis of the tissue factor-dependent coagulation pathway. *J Cell Biol*. 1996;133:293.
- Broze GJ Jr, Lange GW, Duthill KL, MacPhail L. Heterogeneity of plasma tissue factor pathway inhibitor. *Blood Coagul Fibrinolysis*. 1994;5:551.